

predicate logic & dependent types


Freek Wiedijk

Type Theory & Coq

2024–2025

Radboud University Nijmegen

September 27, 2024



introduction

function types become dependent

for predicate logic one needs to generalize

$$(\lambda x : A. M) \text{ : } A \rightarrow B$$

function types

to

$$(\lambda x : A. M) \text{ : } \Pi x : A. B$$

dependent function types

$$\forall x : A. B$$

forall x : A, B

three different notations for the same type

we skip chapters 3 and 5 for now

inductive types



next week

program extraction




last week (briefly)

last week and this week

Curry-Howard correspondence:

propositional logic	$\lambda \rightarrow$	simple types
predicate logic	λP	dependent types
second order logic	$\lambda 2$	polymorphic types
beyond minimal logic	CIC	inductive types

dependent types in the lambda cube



recap: propositional logic

$A, B ::= a \mid A \rightarrow B \mid A \wedge B \mid A \vee B \mid \neg A \mid \top \mid \perp$

$$\begin{array}{ll} I\rightarrow & E\rightarrow \\ I\wedge & El\wedge Er\wedge \\ Il\vee Ir\vee & Ev \\ I\neg & En \\ IT & E\perp \end{array}$$

recap: simply typed lambda calculus

$$\begin{aligned} A, B ::= & a \mid A \rightarrow B \\ M, N ::= & x \mid MN \mid \lambda x : A. M \end{aligned}$$

$$\frac{}{\Gamma \vdash \textcolor{red}{x} : A} \quad \text{for } (x : A) \in \Gamma$$

$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \qquad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash \textcolor{red}{F}M : B}$$

predicate logic

three predicate logics

- ▶ minimal predicate logic

$\rightarrow \forall$

- ▶ constructive predicate logic

$\rightarrow \wedge \vee \neg \top \perp \forall \exists$

- ▶ classical predicate logic

$A \vee \neg A$
 $\neg \neg A \rightarrow A$

propositional versus predicate logic: syntax

$$A, B ::= a \mid A \rightarrow B \mid A \wedge B \mid A \vee B \mid \neg A \mid \top \mid \perp$$
$$M, N ::= x \mid f(\vec{M})$$
$$\vec{M} ::= \cdot \mid \vec{M}, N$$
$$A, B ::= p(\vec{M}) \mid A \rightarrow B \mid A \wedge B \mid A \vee B \mid \neg A \mid \top \mid \perp \mid$$
$$\forall x. A \mid \exists x. A$$

\forall and \exists bind weakly

minimal propositional versus minimal predicate logic: rules

$$\begin{array}{c} [A^H] \\ \vdots \\ \frac{B}{A \rightarrow B} I[H] \rightarrow \end{array} \qquad \qquad \begin{array}{c} \vdots \qquad \vdots \\ \frac{A \rightarrow B \quad A}{B} E \rightarrow \end{array}$$

$$\begin{array}{c} \vdots \\ \frac{A}{\forall x. A} I\forall \end{array} \qquad \qquad \begin{array}{c} \vdots \\ \frac{\forall x. A}{A[x := M]} E\forall \end{array}$$

variable condition of $I\forall$: x not free in available assumptions

same four rules with explicit assumptions

$$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \rightarrow B} I\rightarrow$$

$$\frac{\Gamma \vdash A \rightarrow B \quad \Gamma \vdash A}{\Gamma \vdash B} E\rightarrow$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash \forall x. A} I\forall$$

$$\frac{\Gamma \vdash \forall x. A}{\Gamma \vdash A[x := M]} E\forall$$

variable condition of $I\forall$: x not free in Γ

rules for the existential quantifier

$$\frac{\vdots \quad A[x := M] \quad I\exists}{\exists x. A} \quad \frac{\vdots \quad \exists x. A \quad \forall x. A \rightarrow C \quad E\exists}{C}$$

variable condition of $E\exists$: x not free in C

example proof in minimal logic

$$\frac{\frac{[\forall y. p(y)^H] \quad E\forall}{p(x)} \quad I[H]\rightarrow}{\frac{(\forall y. p(y)) \rightarrow p(x)}{\forall x. ((\forall y. p(y)) \rightarrow p(x))} \quad I\forall}$$

variable condition check:

at the $I\forall$ step the variable x does not occur in assumptions
(there are no available assumptions at that point)

example with existential quantifier

$$A \leftrightarrow B := (A \rightarrow B) \wedge (B \rightarrow A)$$

$$(\exists x. p(x)) \leftrightarrow \neg(\forall x. \neg p(x))$$

not constructively valid

$$\begin{aligned} & (\exists x. p(x)) \rightarrow \neg(\forall x. \neg p(x)) \\ & \neg(\forall x. \neg p(x)) \rightarrow \neg\neg(\exists x. p(x)) \end{aligned}$$

example from left to right

$$\frac{\frac{[\exists x. p(x)^{H_0}] \quad [\forall x. p(x) \rightarrow \perp^{H_1}]}{\perp} E\exists}{\frac{\neg(\forall x. \neg p(x))}{(\exists x. p(x)) \rightarrow \neg(\forall x. \neg p(x))} I[H_0] \rightarrow} I[H_1] \neg$$

variable condition check:

at the $E\exists$ step the variable x does not occur in \perp

example from right to left

$$\frac{\frac{\frac{[\neg(\exists x. p(x))^{H_1}] \quad [p(x)^{H_2}]}{\frac{\frac{\perp}{\neg p(x)} I[H_2]\neg}{\forall x. \neg p(x) E\neg}} I\exists}{\frac{\frac{\perp}{\neg\neg(\exists x. p(x))} I[H_1]\neg}{\forall x. \neg p(x) E\neg}} I\forall}{\neg(\forall x. \neg p(x)) \rightarrow \neg\neg(\exists x. p(x))} I[H_0]\rightarrow$$

variable condition check:

at the $I\forall$ step the variable x does not occur in assumptions

dependent type theory

types thus far

- ▶ atomic types
 a, b, c, \dots
- ▶ types of proof objects of propositions
 a, b, c, \dots
- ▶ function types
- ▶ **datatypes**
 - ▶ natural numbers
 - ▶ Booleans
 - ▶ lists
 - ▶ binary trees
 - ▶ ...

how are types introduced?

- ▶ free type variables

STT = simple type theory

- ▶ in the context = 'axiomatic'

PTSs = pure type systems: $\lambda \rightarrow \lambda P \ \lambda 2 \ \lambda C$

$$\Gamma \vdash M : A$$

$$\text{nat} : *, O : \text{nat}, S : \text{nat} \rightarrow \text{nat} \vdash S(S(O)) : \text{nat}$$

Coq: Parameter

- ▶ definitions → next week

CIC = Calculus of Inductive Constructions

$$E[\Gamma] \vdash M : A$$

Coq: Definition Inductive

star and box

the **sorts** of the lambda cube:

- * = the type of types
- = the type of *

the sort * is a **kind**

= third of four levels

object : type

type : kind

kind : □

object : type : **kind** : □

S : nat → nat : * : □

example contexts

natural numbers:

```
nat : *
O : nat
S : nat → nat
add : nat → nat → nat
```

Booleans:

```
bool : *
true : bool
false : bool
```

lists

lists of Booleans:

list : *

nil : list

cons : bool → list → list

hd : list → bool

tl : list → list

append : list → list → list

reverse : list → list

cons has two arguments
(the head and tail to be put together)

trueslist

the function ‘trueslist’ maps a number n to a list of length n consisting of n copies of ‘true’

trueslist 0 = nil

trueslist (S 0) = cons true nil

trueslist (S (S 0)) = cons true (cons true nil)

trueslist (S (S (S 0))) = cons true (cons true (cons true nil))

...

trueslist : **nat** → list

vectors and truesvec

vectors of Booleans:

$\text{vec} : \text{nat} \rightarrow *$	
$\text{vec } 0 : *$	vectors of length 0
$\text{vec } (\text{S } 0) : *$	vectors of length 1
$\text{vec } (\text{S } (\text{S } 0)) : *$	vectors of length 2
...	

$\text{trueslist} : \text{nat} \rightarrow \text{list}$

$\text{truesvec} : \text{nat} \rightarrow \text{vec } n$

$\text{truesvec} : \prod_{n:\text{nat}} \text{vec } n$

vnil and vcons

`vnil : vec O`

`vcons : nat → bool → vec n → vec (S n)`

`vcons : $\prod_{n : \text{nat}} \text{bool} \rightarrow \text{vec } n \rightarrow \text{vec } (\text{S } n)$`

vcons has three arguments!

`truesvec (S (S O)) : vec (S (S O))`

`truesvec (S (S O)) = vcons (S O) true (vcons O true vnil)`

types when applying vcons

$\text{truesvec } (\mathbf{S} \ O) = \text{vcons } O \ \text{true} \ \text{vnil}$

$\text{truesvec} : \prod n : \text{nat}. \text{vec } n$

$\text{truesvec } (\mathbf{S} \ O) : \text{vec } (\mathbf{S} \ O)$

$\text{vcons} : \prod n : \text{nat}. \text{bool} \rightarrow \text{vec } n \rightarrow \text{vec } (\mathbf{S} \ n)$

$O : \text{nat}$

$\text{vcons } O : \text{bool} \rightarrow \text{vec } O \rightarrow \text{vec } (\mathbf{S} \ O)$

$\text{true} : \text{bool}$

$\text{vcons } O \ \text{true} : \text{vec } O \rightarrow \text{vec } (\mathbf{S} \ O)$

$\text{vnil} : \text{vec } O$

$\text{vcons } O \ \text{true} \ \text{vnil} : \text{vec } (\mathbf{S} \ O)$

BHK-interpretation

proof of $A \rightarrow B$ function from proofs of A to proofs of B

proof of $\neg A$ function from proofs of A to proofs of \perp

proof of $A \wedge B$ pair of a proof of A and a proof of B

proof of $A \vee B$ either a proof of A and a proof of B

proof of $\forall x. A$ dependent function from objects to proofs of A

proof of $\exists x. A$ dependent pair of an object and a proof of A

dependent = A is not fixed but *depends* on the object x

$$A \vee B \leftrightarrow \exists x \in \{\text{left, right}\}. (x = \text{left} \rightarrow A) \wedge (x = \text{right} \rightarrow B)$$

Curry-Howard correspondence

type theory	Coq	minimal logic
$A \rightarrow B$	$A \rightarrow B$	$A \rightarrow B$
$\Pi x : D. A$	<code>forall x : D, A</code>	$\forall x. A$

$D : *$ is called Terms in the notes

proof term	type	statement proved
$H : A \rightarrow B$	$A \rightarrow B$	$A \rightarrow B$
$H' : A$		A
$HH' : B$		B
$H : \Pi x : D. Px$		$\forall x. P(x)$
$HM : PM$		$P(M)$

Coq version of the first example

```
Parameter D : Set.          four =
Parameter p : D -> Prop.    fun (x : D) (H : forall y : D, p y)
                            => H x
Lemma four :                  : forall x : D,
forall x : D,                      (forall y : D, p y) -> p x
  (forall y : D, p y) ->
  p x.                                Arguments four _ _%function_scope
intros x H.                           Qed.
```

Print four.

$$\frac{\frac{[\forall y. p(y)^H]}{p(x)} E\forall}{(\forall y. p(y)) \rightarrow p(x)} I[H]\rightarrow$$
$$\frac{(\forall y. p(y)) \rightarrow p(x)}{\forall x. (\forall y. p(y)) \rightarrow p(x)} I\forall$$

$$(\lambda x : D. \lambda H : (\Pi y : D. py). Hx) : \Pi x : D. (\Pi y : D. py) \rightarrow px$$

understanding the proof term

$$\forall x. (\forall y. p(y)) \rightarrow p(x)$$

$$\lambda x : D. \lambda H : (\Pi y : D. py). Hx$$

the second example in Coq, from left to right

```
Parameter D : Set.  
Parameter p : D -> Prop.  
  
Lemma five :  
  (exists x : D, p x) ->  
  ~ (forall x : D, ~ p x).  
intros H0 H1.  
elim H0.  
apply H1.  
Qed.
```

$$\frac{\frac{[\exists x. p(x)^{H_0}] \quad [\forall x. p(x) \rightarrow \perp^{H_1}]}{\perp} E\exists}{\neg(\forall x. \neg p(x)) I[H_1]\neg} I[H_0]\rightarrow$$

the second example in Coq, from right to left

```
Parameter D : Set.
```

```
Parameter p : D -> Prop.
```

Lemma six :

```
~ (forall x : D, ~ p x) ->
~ ~ (exists x : D, p x).
```

```
intros H0 H1.
```

```
apply H0.
```

```
intros x H2.
```

```
apply H1.
```

```
exists x.
```

```
apply H2.
```

```
Qed.
```

$$\frac{\frac{\frac{\frac{[\neg(\exists x. p(x))^{H_1}] \quad [p(x)^{H_2}]}{\exists x. p(x)} I_{\exists}}{\perp} E_{\rightarrow}}{\neg p(x)} I[H_2] \neg}{\frac{\frac{[\neg(\forall x. \neg p(x))^{H_0}] \quad [\forall x. \neg p(x)]}{\perp} I_{\forall}}{\neg \neg(\exists x. p(x))} E_{\neg}} I[H_1] \neg}{\neg(\forall x. \neg p(x)) \rightarrow \neg \neg(\exists x. p(x))} I[H_0] \rightarrow$$

proof rules versus Coq tactics

$I \rightarrow I\forall$	intro intros
$E \rightarrow E\forall$	apply

$E \wedge E\vee E\exists$	elim destruct intro-patterns
$I \wedge$	split
$I \vee$	left right
$I \exists$	exists

detours

proof normalization

detour = introduction rule directly followed by a elimination rule for the same connective

cut = corresponding notion in sequent calculus

can be eliminated → reduction of the proof term

detour for implication:

'prove a lemma A and then prove B using this lemma'

detour elimination is inlining the proof₁ of the lemma everywhere

$$\frac{\frac{[A^H]}{\vdots_2} \frac{\frac{B}{A \rightarrow B} I[H] \rightarrow \frac{\vdots_1}{A}}{\frac{E \rightarrow}{B}} \longrightarrow_{\beta}}{\vdots_1 A} \vdots_2 B$$

detours for predicate logic

detour for the universal quantifier

generalize the statement $A[x := M]$ to A with arbitrary x
elimination is specializing the proof₁ to M

$$\frac{\vdots_1 \quad A}{\forall x. A} I\forall \quad \frac{A[x := M]}{A[x := M]} E\forall \quad \rightarrow_{\beta} \quad \frac{\vdots_1[x := M]}{A[x := M]}$$

$$(\lambda x : D. H_1)M \rightarrow_{\beta} H_1[x := M]$$

STT versus $\lambda\mathbf{P}$ syntax

STT:

$$\begin{aligned} A, B ::= & \textcolor{red}{a} \mid A \rightarrow B \\ M, N ::= & \textcolor{red}{x} \mid MN \mid \lambda x : A. M \\ \Gamma ::= & \cdot \mid \Gamma, x : A \\ \mathcal{J} ::= & \Gamma \vdash M : A \end{aligned}$$

$\lambda\mathbf{P}$:

$$\begin{aligned} s ::= & * \mid \square \\ M, N, A, B ::= & \textcolor{red}{x} \mid MN \mid \lambda x : A. M \mid \frac{}{\Pi x : A. B} \mid s \end{aligned}$$

$$\begin{aligned} \Gamma ::= & \cdot \mid \Gamma, x : A \\ \mathcal{J} ::= & \Gamma \vdash M : A \end{aligned}$$

$$\frac{\Gamma}{a : *, b : *, f : a \rightarrow b, x : a \vdash fx : b} \frac{M \quad A}{\Pi x : A. B}$$

STT versus λ P

STT

3 rules

$$x \mid MN \mid \lambda x : A. M$$

λ P

7 rules

$$x \mid MN \mid \lambda x : A. M \mid \Pi x : A. M \mid *$$

box does not have a type
→ no rule for box

variables have two rules

conversion rule

typing rules

STT

λP

$$\frac{}{\vdash * : \square}$$

axiom rule
start rule

$$\frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : s}{\Gamma \vdash \Pi x : A. B : s}$$

product rule

$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x : A. M : A \rightarrow B}$$

$$\frac{\Gamma, x : A \vdash M : B \quad \Gamma \vdash \Pi x : A. B : s}{\Gamma \vdash \lambda x : A. M : \Pi x : A. B}$$

abstraction rule

typing rules (continued)

STT

λP

$$\frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

$$\frac{\Gamma \vdash F : \Pi x : A. B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B[x := M]}$$

application rule

$$\begin{aligned} \text{truesvec} &: \Pi n : \text{nat}. \text{vec } n \\ \text{truesvec } (\text{S } (\text{S } O)) &: \text{vec } (\text{S } (\text{S } O)) \end{aligned}$$

typing rules (continued)

STT

(context Γ is a set)

λP

(context Γ is a list)

correct: $a : *, x : a \vdash x : a$

incorrect: $x : a, a : * \vdash x : a$

$$\frac{}{\Gamma \vdash x : A} (x : A) \in \Gamma$$

$$\frac{\Gamma \vdash A : s}{\Gamma, x : A \vdash x : A}$$

variable rule

$$\frac{\Gamma \vdash M : A \quad \Gamma \vdash B : s}{\Gamma, y : B \vdash M : A}$$

weakening rule

the final typing rule

$\text{vecappend} : \Pi n_1 : \text{nat}. \Pi n_2 : \text{nat}.$

$\text{vec } n_1 \rightarrow \text{vec } n_2 \rightarrow \text{vec } (\text{add } n_1 n_2)$

$\text{vecappend } 3\ 4\ v_1\ v_2 : \text{vec } (\text{add } 3\ 4)$

$\text{vecappend } 3\ 4\ v_1\ v_2 : \text{vec } 7$

λP

$$\frac{\Gamma \vdash M : A \quad \Gamma \vdash A' : s}{\Gamma \vdash M : A'} A =_{\beta} A'$$

conversion rule

A and A' are convertible

$=_{\beta}$ is defined on preterms

λP on one slide

$$\begin{aligned} M, F, A, B ::= & \textcolor{red}{x} \mid \textcolor{red}{F}M \mid \lambda x : A. M \mid \Pi x : A. B \mid s \\ s ::= & * \mid \square \end{aligned}$$

$$\frac{}{\vdash * : \square} \quad \frac{\Gamma \vdash A : s}{\Gamma, x : A \vdash \textcolor{red}{x} : A} \quad \frac{\Gamma \vdash M : A \quad \Gamma \vdash B : s}{\Gamma, \textcolor{green}{y} : \textcolor{blue}{B} \vdash M : A}$$

$$\frac{\Gamma, x : A \vdash M : B \quad \Gamma \vdash \Pi x : A. B : s}{\Gamma \vdash \lambda x : A. M : \Pi x : A. B} \quad \frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : s}{\Gamma \vdash \Pi x : A. B : s}$$

$$\frac{\Gamma \vdash F : \Pi x : A. B \quad \Gamma \vdash M : A}{\Gamma \vdash \textcolor{red}{F}M : B[x := M]} \quad \frac{\Gamma \vdash M : A \quad \Gamma \vdash A' : s}{\Gamma \vdash M : \textcolor{green}{A}'} A =_{\beta} A'$$

the lambda cube


only the product rules differ:

$$\lambda\rightarrow \frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : *}{\Gamma \vdash \Pi x : A. B : *}$$

$$\lambda P \frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : \textcolor{red}{s}}{\Gamma \vdash \Pi x : A. B : \textcolor{red}{s}}$$

$$\lambda 2 \frac{\Gamma \vdash A : \textcolor{red}{s} \quad \Gamma, x : A \vdash B : *}{\Gamma \vdash \Pi x : A. B : *}$$

$$\lambda C \frac{\Gamma \vdash A : \textcolor{red}{s}_1 \quad \Gamma, x : A \vdash B : \textcolor{red}{s}_2}{\Gamma \vdash \Pi x : A. B : \textcolor{red}{s}_2}$$



pure type systems

$$\frac{}{\vdash s_1 : s_2} (s_1, s_2) \in \mathcal{A}$$

$$\frac{\Gamma \vdash A : s_1 \quad \Gamma, x : A \vdash B : s_2}{\Gamma \vdash \Pi x : A. B : s_3} (s_1, s_2, s_3) \in \mathcal{R}$$

	\mathcal{S} sorts	\mathcal{A} axioms	\mathcal{R} rules
$\lambda*$	$\{*\}$	$\{(*, *)\}$	$\{(*, *, *)\}$
$\lambda\rightarrow$	$\{*, \square\}$	$\{(*, \square)\}$	$\{(*, *, *)\}$
λP	$\{*, \square\}$	$\{(*, \square)\}$	$\{(*, *, *), (*, \square, \square)\}$
λC	$\{*, \square\}$	$\{(*, \square)\}$	$\{(*, *, *), (\ast, \square, \square),$ $(\square, *, *), (\square, \square, \square)\}$
$\lambda PRED$	$\{*_s, \square_s, *_p, \square_p\}$	$\{(*_s, \square_s), (*_p, \square_p)\}$	$\{(*_s, *_s, *_s), (*_s, \square_s, \square_s),$ $(*_p, *_p, *_p), (*_s, *_p, *_p)\}$

example PTS type derivation

$$\frac{\frac{}{\vdash * : \square} \quad \frac{}{\vdash * : \square}}{a : * \vdash a : * \quad a : * \vdash a : *} \quad \text{(next slide)} \quad \vdots \quad \frac{\vdash * : \square}{a : * \vdash a : *}$$
$$\frac{a : *, x : a \vdash a : * \quad \frac{a : *, x : a, y : a \vdash y : a \quad a : *, x : a \vdash a \rightarrow a : *}{a : *, x : a \vdash (\lambda y : a. y) : a \rightarrow a} \quad a : *, x : a \vdash x : a}{a : *, x : a \vdash (\lambda y : a. y)x : a}$$

$$\frac{\frac{x : a, y : a \vdash y : a}{x : a \vdash (\lambda y : a. y) : a \rightarrow a} \quad \frac{}{x : a \vdash x : a}}{x : a \vdash (\lambda y : a. y)x : a}$$

example PTS type derivation (continued)

$$\frac{\frac{\frac{\frac{\frac{\frac{\frac{}{\vdash * : \square}}{\vdash * : \square} \quad \frac{\frac{\frac{\vdash * : \square}{\vdash * : \square}}{a : * \vdash a : *} \quad \frac{\vdash * : \square}{a : * \vdash a : *}}{a : * \vdash a : *} \quad \frac{}{\vdash * : \square}}{a : * \vdash a : *} \quad \frac{a : *, y : a \vdash a : *}{a : *, y : a \vdash a : *}}{a : * \vdash (\Pi y : a. a) : *} \quad \frac{}{a : * \vdash a : *}$$
$$\frac{a : *, x : a \vdash \underline{a \rightarrow a} : *}{\Pi y : a. a}$$

$$\frac{}{\vdash * : \square} \quad \frac{\Gamma \vdash A : s}{\Gamma, x : A \vdash x : A} \quad \frac{\Gamma \vdash M : A \quad \Gamma \vdash B : s}{\Gamma, y : B \vdash M : A}$$

$$\frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : s}{\Gamma \vdash \Pi x : A. B : s}$$

conclusion

computing the sort of a type

$$\text{type}_\Gamma(\lambda x : A. M) = \Pi x : A. \text{type}_{\Gamma, x:A}(M)$$

$$\text{type}_\Gamma(\Pi x : A. B) = \text{type}_{\Gamma, x:A}(B)$$

$$\text{type}_\Gamma(A \rightarrow B) = \text{type}_\Gamma(B)$$

$$\text{type}_\Gamma(*) = \square$$

$$\text{type}_\Gamma(x) = \Gamma(x)$$

$$\text{type}_\Gamma(FM) = \dots$$

summary

- ▶ predicate logic
- ▶ dependent types
 - ▶ vectors
 - ▶ Curry-Howard
- ▶ λP
 - ▶ lambda cube
 - ▶ PTSs
- ▶ detours

table of contents

contents

introduction

predicate logic

dependent type theory

Coq

detours

λP

conclusion

table of contents