

propositional logic & simple types


Freek Wiedijk

Type Theory & Rocq

2025–2026

Radboud University Nijmegen

September 10, 2025



type systems and logics

teaser: a proof term for a proof

$\lambda x : a \rightarrow b \rightarrow c. \lambda y : a \rightarrow b. \lambda z : a. xz(yz)$

$$\frac{\frac{\frac{[a \rightarrow b \rightarrow c^x] \quad [a^z]}{b \rightarrow c} E \rightarrow \quad \frac{\frac{[a \rightarrow b^y] \quad [a^z]}{b} E \rightarrow}{c} E \rightarrow}{\frac{c}{a \rightarrow c} I[z] \rightarrow} I[y] \rightarrow \frac{(a \rightarrow b) \rightarrow a \rightarrow c}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow$$

many systems

- ▶ one untyped lambda calculus
(last week: recap)
- ▶ many typed lambda calculi = type theories
 - ▶ **Church-style**
variables explicitly typed

the systems in this course
Rocq!
 - ▶ **Curry-style**
assigning types to untyped terms
(Herman's lecture next week)
- ▶ many logics

Curry-Howard correspondence

'propositions-as-types'

type systems $\sim$ logics

datatypes $\sim$ propositions

$A \times B$ $\sim$ $A \wedge B$

$A \rightarrow B$ $\sim$ $A \rightarrow B$

objects of a datatype $\sim$ proofs of a proposition
'inhabitants' $\sim$ 'proof objects'

non-empty type $\sim$ true proposition

empty type $\sim$ false proposition

Curry-Howard correspondence

'propositions-as-types'

type systems $\sim$ logics

datatypes $\sim$ propositions

$A \times B$ $\sim$ $A \wedge B$

$A \rightarrow B$ $\sim$ $A \rightarrow B$

objects of a datatype $\sim$ proofs of a proposition
'inhabitants' $\sim$ 'proof objects'

non-empty type $\sim$ true proposition

empty type $\sim$ false proposition

derivation in a type theory $\sim$ derivation in a logic

type systems in this course

logics $\sim$ type systems


propositional logic $\sim \lambda \rightarrow$ = simply typed lambda calculus

predicate logic $\sim \lambda P$ = dependently typed lambda calculus

second order logic $\sim \lambda 2$ = polymorphic lambda calculus

higher order logic $\sim \lambda C$ = CC = Calculus of Constructions

'the logic of Rocq' $\sim$ CIC = Calculus of Inductive Constructions




the lambda cube

the Barendregt cube

eight PTSs = pure type systems

(not explicitly in Femke's course notes)



► **natural deduction**

$$B_1, \dots, B_m \vdash A$$

introduction and elimination rules for each connective

$$\frac{\dots \vdash \dots}{\dots \vdash \dots \otimes \dots} I$$

$$\frac{\dots \vdash \dots \otimes \dots}{\dots \vdash \dots} E$$

- Gentzen–Prawitz style: proof is a derivation tree
- Jaśkowski–Fitch style: proof consists of nested boxes/flags

► **sequent calculus**: LK & LJ

$$B_1, \dots, B_m \vdash A_1, \dots, A_n$$

left and right rules for each connective

$$\frac{\dots \vdash \dots}{\dots \otimes \dots \vdash \dots} L$$

$$\frac{\dots \vdash \dots}{\dots \vdash \dots \otimes \dots} R$$

► **Hilbert-style**

$$\vdash A$$

axioms for each connective (and at most two rules)

► **natural deduction**

$$B_1, \dots, B_m \vdash A$$

introduction and elimination rules for each connective

$$\frac{\dots \vdash \dots}{\dots \vdash \dots \otimes \dots} I$$

$$\frac{\dots \vdash \dots \otimes \dots}{\dots \vdash \dots} E$$

- **Gentzen–Prawitz style:** proof is a derivation tree
- **Jaśkowski–Fitch style:** proof consists of nested boxes/flags

► **sequent calculus:** LK & LJ

$$B_1, \dots, B_m \vdash A_1, \dots, A_n$$

left and right rules for each connective

$$\frac{\dots \vdash \dots}{\dots \otimes \dots \vdash \dots} L$$

$$\frac{\dots \vdash \dots}{\dots \vdash \dots \otimes \dots} R$$

► **Hilbert-style**

$$\vdash A$$

axioms for each connective (and at most two rules)

defining a logic or type system

- ▶ **syntax**

- ▶ terms
- ▶ types
- ▶ formulas
- ▶ contexts
- ▶ judgments

- ▶ **rules**

- ▶ **logics**: proof rules
- ▶ **type systems**: typing rules

- ▶ **reduction**

defining a logic or type system

► syntax

- terms
- types
- formulas
- contexts
- judgments

$$M, N ::= x \mid MN \mid \lambda x. M$$

► rules

- **logics**: proof rules
- **type systems**: typing rules

no rules for untyped lambda calculus

► reduction

$$(\lambda x. M) N \rightarrow_{\beta} M[x := N]$$

propositional logic

three propositional logics

- ▶ **minimal propositional logic**

the logic that corresponds to simply typed lambda calculus

only implication: $\rightarrow$

- ▶ **constructive propositional logic**

all connectives: $\rightarrow, \wedge, \vee, \neg, \perp, \top$

- ▶ **classical propositional logic**

$$\begin{aligned} & A \vee \neg A \\ & \neg\neg A \rightarrow A \end{aligned}$$

propositional logic: syntax

formulas

$$A, B ::= a \mid A \rightarrow B \mid A \wedge B \mid A \vee B \mid \neg A \mid \top \mid \perp$$

$a, b, c, \dots$ atomic propositions
later: propositional variables

$A, B, C, \dots$ meta-variables for arbitrary formulas

implication associates to the right:

$$a \rightarrow b \rightarrow c \text{ means } a \rightarrow (b \rightarrow c)$$

order of binding strength: $\neg > \wedge > \vee > \rightarrow$

$$a \vee \neg b \wedge c \rightarrow d \text{ means } (a \vee ((\neg b) \wedge c)) \rightarrow d$$

the two rules of minimal propositional logic

the introduction and elimination rules of implication:

$$\frac{\begin{array}{c} [A^x] \\ \vdots \\ B \end{array}}{A \rightarrow B} I[x] \rightarrow \qquad \frac{\begin{array}{c} \vdots \\ A \rightarrow B \end{array} \quad \begin{array}{c} \vdots \\ A \end{array}}{B} E \rightarrow$$

in the introduction rule the assumption $[A^x]$ may be used an arbitrary number of times: zero, one or more

in the elimination rule the proof of $A \rightarrow B$ is on the *left* of the proof of A

example proof of minimal propositional logic

$$(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c$$

example proof of minimal propositional logic

$[a \rightarrow b \rightarrow c^x]$

$$\frac{(a \rightarrow b) \rightarrow a \rightarrow c}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y]$$

$$\frac{\frac{a \rightarrow c}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y] \rightarrow}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$$

$$\frac{\frac{\frac{c}{a \rightarrow c} I[z] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y] \rightarrow}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$$

$$\frac{\frac{\frac{b \rightarrow c}{a \rightarrow c} I[z] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y] \rightarrow}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow \quad \frac{b}{E \rightarrow}$$

example proof of minimal propositional logic

$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$

$$\frac{\frac{a \rightarrow b \rightarrow c \quad a}{b \rightarrow c} E \rightarrow \quad b}{\frac{\frac{\frac{c}{a \rightarrow c} I[z] \rightarrow \quad (a \rightarrow b) \rightarrow a \rightarrow c}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y] \rightarrow \quad (a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow} E \rightarrow$$

example proof of minimal propositional logic

$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$

$$\begin{array}{c}
 \frac{[a \rightarrow b \rightarrow c^x] \quad a}{b \rightarrow c} E\rightarrow \\
 \frac{\frac{\frac{b}{a \rightarrow b} E\rightarrow}{c} I[z]\rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y]\rightarrow \\
 \frac{(a \rightarrow b) \rightarrow a \rightarrow c}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x]\rightarrow
 \end{array}$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$$

$$\begin{array}{c}
 \frac{[a \rightarrow b \rightarrow c^x] \quad [a^z]}{b \rightarrow c} E \rightarrow \qquad b \\
 \hline
 \qquad \qquad \qquad \frac{b}{E \rightarrow} \\
 \qquad \qquad \qquad \frac{c}{a \rightarrow c} I[z] \rightarrow \\
 \qquad \qquad \qquad \frac{a \rightarrow c}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y] \rightarrow \\
 \hline
 (a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c \quad I[x] \rightarrow
 \end{array}$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$$

$$\frac{\frac{[a \rightarrow b \rightarrow c^x] \quad [a^z]}{b \rightarrow c} E\rightarrow \quad \frac{a \rightarrow b \quad a}{b} E\rightarrow}{\frac{c}{a \rightarrow c} I[z]\rightarrow} E\rightarrow$$

$$\frac{\frac{a \rightarrow c}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y]\rightarrow}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x]\rightarrow$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$$

$$\begin{array}{c}
 \frac{[a \rightarrow b \rightarrow c^x] \quad [a^z]}{b \rightarrow c} E\rightarrow \quad \frac{[a \rightarrow b^y] \quad a}{b} E\rightarrow \\
 \hline
 \frac{\frac{c}{a \rightarrow c} I[z]\rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y]\rightarrow \\
 \hline
 (a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c \quad I[x]\rightarrow
 \end{array}$$

example proof of minimal propositional logic

$$[a \rightarrow b \rightarrow c^x] \quad [a \rightarrow b^y] \quad [a^z]$$

$$\frac{\frac{[a \rightarrow b \rightarrow c^x] \quad [a^z]}{b \rightarrow c} E \rightarrow \quad \frac{[a \rightarrow b^y] \quad [a^z]}{b} E \rightarrow}{\frac{c}{a \rightarrow c} I[z] \rightarrow} E \rightarrow$$

$$\frac{\frac{a \rightarrow c}{(a \rightarrow b) \rightarrow a \rightarrow c} I[y] \rightarrow}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow$$

example proof of minimal propositional logic

$$\frac{\frac{\frac{[a \rightarrow b \rightarrow c^x] \quad [a^z]}{b \rightarrow c} E \rightarrow \quad \frac{\frac{[a \rightarrow b^y] \quad [a^z]}{b} E \rightarrow}{b} E \rightarrow}{\frac{c}{a \rightarrow c} I[z] \rightarrow} I[y] \rightarrow \frac{(a \rightarrow b) \rightarrow a \rightarrow c}{(a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I[x] \rightarrow$$

constructive propositional logic: the other rules

$$\frac{\begin{array}{c} \vdots \\ A \end{array} \quad \begin{array}{c} \vdots \\ B \end{array}}{A \wedge B} I\wedge \quad \frac{\begin{array}{c} \vdots \\ A \wedge B \end{array}}{A} El\wedge \quad \frac{\begin{array}{c} \vdots \\ A \wedge B \end{array}}{B} Er\wedge$$

$$\frac{\begin{array}{c} \vdots \\ A \end{array}}{A \vee B} Il\vee \quad \frac{\begin{array}{c} \vdots \\ B \end{array}}{A \vee B} Ir\vee \quad \frac{\begin{array}{c} \vdots \\ A \vee B \end{array} \quad \begin{array}{c} \vdots \\ A \rightarrow C \end{array} \quad \begin{array}{c} \vdots \\ B \rightarrow C \end{array}}{C} EV$$

$$\frac{\begin{array}{c} [A^x] \\ \vdots \\ \perp \end{array}}{\neg A} I\neg \quad \frac{\begin{array}{c} \vdots \\ \neg A \end{array} \quad \begin{array}{c} \vdots \\ A \end{array}}{\perp} E\neg \quad \frac{}{\top} I\top \quad \frac{\perp}{A} E\perp$$

constructive propositional logic: the other rules

$$\begin{array}{c}
 \begin{array}{c} \vdots \\ A \end{array} \quad \begin{array}{c} \vdots \\ B \end{array} \\
 \hline
 A \wedge B \quad I\wedge
 \end{array}
 \quad
 \begin{array}{c}
 \begin{array}{c} \vdots \\ A \wedge B \end{array} \\
 \hline
 A \quad El\wedge
 \end{array}
 \quad
 \begin{array}{c}
 \begin{array}{c} \vdots \\ A \wedge B \end{array} \\
 \hline
 B \quad Er\wedge
 \end{array}$$

$$\begin{array}{c}
 \begin{array}{c} \vdots \\ A \end{array} \\
 \hline
 A \vee B \quad Il\vee
 \end{array}
 \quad
 \begin{array}{c}
 \begin{array}{c} \vdots \\ B \end{array} \\
 \hline
 A \vee B \quad Ir\vee
 \end{array}
 \quad
 \begin{array}{c}
 \begin{array}{c} \vdots \\ A \vee B \end{array} \quad \begin{array}{c} \vdots \\ A \rightarrow C \end{array} \quad \begin{array}{c} \vdots \\ B \rightarrow C \end{array} \\
 \hline
 C \quad E\vee
 \end{array}$$

$$\begin{array}{c}
 [A^x] \\
 \begin{array}{c} \vdots \\ \perp \end{array} \\
 \hline
 \neg A \quad I\neg
 \end{array}
 \quad
 \begin{array}{c}
 \begin{array}{c} \vdots \\ \neg A \end{array} \quad \begin{array}{c} \vdots \\ A \end{array} \\
 \hline
 \perp \quad E\neg
 \end{array}
 \quad
 \begin{array}{c}
 \overline{\top} \quad I\top
 \end{array}
 \quad
 \begin{array}{c}
 \begin{array}{c} \vdots \\ \perp \end{array} \\
 \hline
 A \quad E\perp
 \end{array}$$

$$\neg A := A \rightarrow \perp$$

variants of elimination rules

- often the disjunction elimination rule is:

$$\frac{\begin{array}{ccc} & [A^x] & [B^y] \\ & \vdots & \vdots \\ A \vee B & C & C \end{array}}{C} E[x, y] \vee$$

not what Rocq's `elim` tactic implements for disjunction

variants of elimination rules

- often the disjunction elimination rule is:

$$\frac{\begin{array}{ccc} & [A^x] & [B^y] \\ \vdots & \vdots & \vdots \\ A \vee B & C & C \end{array}}{C} E[x, y] \vee$$

not what Rocq's `elim` tactic implements for disjunction

- what Rocq's `elim` tactic implements for conjunction:

$$\frac{\begin{array}{cc} \vdots & \vdots \\ A \wedge B & A \rightarrow B \rightarrow C \end{array}}{C} E \wedge$$

example proof beyond minimal propositional logic

$$a \vee b \rightarrow b \vee a$$

example proof beyond minimal propositional logic

$[a \vee b^x]$

$$\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow$$

example proof beyond minimal propositional logic

$[a \vee b^x]$

$$\frac{[a \vee b^x] \quad a \rightarrow b \vee a \quad b \rightarrow b \vee a}{b \vee a} E\vee$$
$$\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow$$

example proof beyond minimal propositional logic

$[a \vee b^x] \quad [a^y]$

$$\frac{\begin{array}{c} \frac{b \vee a}{a \rightarrow b \vee a} I[y] \rightarrow \\ [a \vee b^x] \end{array} \quad \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow}{b \rightarrow b \vee a} E\vee$$

example proof beyond minimal propositional logic

$[a \vee b^x] \quad [a^y]$

$$\begin{array}{c}
 \frac{a}{b \vee a} Ir\vee \\
 \frac{\quad}{a \rightarrow b \vee a} I[y] \rightarrow \\
 \frac{[a \vee b^x] \quad \frac{\quad}{a \rightarrow b \vee a} I[y] \rightarrow \quad b \rightarrow b \vee a}{\quad} E\vee \\
 \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow
 \end{array}$$

example proof beyond minimal propositional logic

$[a \vee b^x] \quad [a^y]$

$$\begin{array}{c}
 \frac{[a^y]}{b \vee a} Ir\vee \\
 \frac{[a \vee b^x] \quad \frac{b \vee a}{a \rightarrow b \vee a} I[y] \rightarrow \quad b \rightarrow b \vee a}{b \vee a} E\vee \\
 \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow
 \end{array}$$

example proof beyond minimal propositional logic

$[a \vee b^x]$

$[b^z]$

$$\begin{array}{c}
 \frac{\frac{\frac{[a^y]}{b \vee a} Ir\vee}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{b \vee a}{b \rightarrow b \vee a} I[z] \rightarrow}{[a \vee b^x] \quad \frac{b \vee a}{a \vee b \rightarrow b \vee a} E\vee} E\vee \\
 \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow
 \end{array}$$

example proof beyond minimal propositional logic

$[a \vee b^x]$

$[b^z]$

$$\begin{array}{c}
 \frac{\frac{\frac{[a^y]}{b \vee a} Ir\vee}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{\frac{b}{b \vee a} Il\vee}{b \rightarrow b \vee a} I[z] \rightarrow}{[a \vee b^x] \quad \frac{b \vee a}{a \vee b \rightarrow b \vee a} E\vee} I[x] \rightarrow \\
 \textcolor{red}{a \vee b \rightarrow b \vee a}
 \end{array}$$

example proof beyond minimal propositional logic

$[a \vee b^x]$

$[b^z]$

$$\begin{array}{c}
 \frac{\frac{\frac{[a^y]}{b \vee a} Ir\vee}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{\frac{[b^z]}{b \vee a} Il\vee}{b \rightarrow b \vee a} I[z] \rightarrow}{[a \vee b^x] \quad \quad \quad E\vee} \\
 \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow
 \end{array}$$

example proof beyond minimal propositional logic

$$\begin{array}{c}
 \frac{[a \vee b^x] \quad \frac{\frac{\frac{[a^y]}{b \vee a} Ir\vee}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{\frac{\frac{[b^z]}{b \vee a} Il\vee}{b \rightarrow b \vee a} I[z] \rightarrow}{E\vee}}{b \vee a} \\
 \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow
 \end{array}$$

alternative style: explicit assumption lists

syntax

$A, B ::= a \mid A \rightarrow B \mid \dots$	formulas
$\Gamma ::= \cdot \mid \Gamma, A$	assumption lists
$\mathcal{J} ::= \Gamma \vdash A$	sequents

we do not write the dot and the comma after the dot
still natural deduction, *not* sequent calculus

rules

$$\frac{}{\Gamma \vdash A} \text{ass} \quad \text{for } A \in \Gamma$$
$$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \rightarrow B} I \rightarrow \quad \frac{\Gamma \vdash A \rightarrow B \quad \Gamma \vdash A}{\Gamma \vdash B} E \rightarrow \quad \dots$$

the example proofs with explicit assumption lists

$\Gamma_1 := a \rightarrow b \rightarrow c, a \rightarrow b, a$

$$\begin{array}{c}
 \frac{\frac{\frac{}{\Gamma_1 \vdash a \rightarrow b \rightarrow c} \text{ass}}{\Gamma_1 \vdash b \rightarrow c} E \rightarrow \quad \frac{\frac{\frac{}{\Gamma_1 \vdash a} \text{ass}}{\Gamma_1 \vdash b} E \rightarrow \quad \frac{\frac{\frac{}{\Gamma_1 \vdash a \rightarrow b} \text{ass}}{\Gamma_1 \vdash a} \text{ass}}{\Gamma_1 \vdash b} E \rightarrow}{\Gamma_1 \vdash c} E \rightarrow \\
 \frac{\Gamma_1 \vdash c}{a \rightarrow b \rightarrow c, a \rightarrow b \vdash a \rightarrow c} I \rightarrow \\
 \frac{a \rightarrow b \rightarrow c, a \rightarrow b \vdash a \rightarrow c}{a \rightarrow b \rightarrow c \vdash (a \rightarrow b) \rightarrow a \rightarrow c} I \rightarrow \\
 \frac{a \rightarrow b \rightarrow c \vdash (a \rightarrow b) \rightarrow a \rightarrow c}{\vdash (a \rightarrow b \rightarrow c) \rightarrow (a \rightarrow b) \rightarrow a \rightarrow c} I \rightarrow \\
 \\
 \frac{\frac{\frac{\frac{}{a \vee b, a \vdash a} \text{ass}}{a \vee b, a \vdash b \vee a} Ir \vee \quad \frac{\frac{\frac{}{a \vee b, b \vdash b} \text{ass}}{a \vee b, b \vdash b \vee a} Il \vee}{a \vee b \vdash a \rightarrow b \vee a} I \rightarrow \quad \frac{\frac{\frac{}{a \vee b, b \vdash b} \text{ass}}{a \vee b, b \vdash b \vee a} Il \vee}{a \vee b \vdash b \rightarrow b \vee a} I \rightarrow}{a \vee b \vdash a \vee b} \text{ass} \quad \frac{a \vee b \vdash a \rightarrow b \vee a \quad a \vee b \vdash b \rightarrow b \vee a}{a \vee b \vdash b \vee a} E \vee}{\vdash a \vee b \rightarrow b \vee a} I \rightarrow
 \end{array}$$

simply typed lambda calculus

the S combinator

$$\lambda xyz. xz(yz)$$

simply typed lambda calculus

the S combinator

$$\lambda xyz. xz(yz)$$
$$(\lambda x. (\lambda y. (\lambda z. ((xz)(yz)))))$$

simply typed lambda calculus

the S combinator

untyped

typed: Curry-style

$$\lambda xyz. xz(yz)$$
$$(\lambda x. (\lambda y. (\lambda z. ((xz)(yz))))))$$

typed: Church-style

$$\lambda x : a \rightarrow b \rightarrow c. \lambda y : a \rightarrow b. \lambda z : a. xz(yz)$$

simply typed lambda calculus

the S combinator

untyped

typed: Curry-style

$$\lambda xyz. xz(yz)$$
$$(\lambda x. (\lambda y. (\lambda z. ((xz)(yz)))))$$

typed: Church-style

$$\lambda x : a \rightarrow b \rightarrow c. \lambda y : a \rightarrow b. \lambda z : a. xz(yz)$$

Rocq syntax

```
fun x : a -> b -> c => fun y : a -> b => fun z : a =>
  x z (y z)
```

simply typed lambda calculus

the S combinator

untyped

typed: Curry-style

$$\lambda xyz. xz(yz)$$
$$(\lambda x. (\lambda y. (\lambda z. ((xz)(yz)))))$$

typed: Church-style

$$\lambda x : a \rightarrow b \rightarrow c. \lambda y : a \rightarrow b. \lambda z : a. xz(yz)$$

Rocq syntax

```
fun x : a -> b -> c => fun y : a -> b => fun z : a =>
  x z (y z)
```

```
fun (x : a -> b -> c) (y : a -> b) (z : a) => x z (y z)
```

BHK interpretation

Brouwer–Heyting–Kolmogorov

~ Curry-Howard correspondence

constructive ‘meaning’ of the logical connectives

connection between proofs and lambda calculus

proof of $A \rightarrow B$	=	function from proofs of A to proofs of B
proof of $A \wedge B$	=	pair of a proof of A and a proof of B
proof of $A \vee B$	=	either a proof of A , or a proof of B
proof of $\neg A$	=	proof of $A \rightarrow \perp$
proof of $\top$		the object I
proof of $\perp$		does not exist

BHK interpretation

Brouwer–Heyting–Kolmogorov

~ Curry-Howard correspondence

constructive ‘meaning’ of the logical connectives

connection between proofs and lambda calculus

proof of $A \rightarrow B$	=	function from proofs of A to proofs of B
proof of $A \wedge B$	=	pair of a proof of A and a proof of B
proof of $A \vee B$	=	either a proof of A , or a proof of B
proof of $\neg A$	=	proof of $A \rightarrow \perp$
proof of $\top$		the object I
proof of $\perp$		does not exist

proof of $a \wedge b \rightarrow b \wedge a$ is

a function that inputs a pair $\langle x, y \rangle$, and returns $\langle y, x \rangle$
with x a proof of a and y a proof of b

$$\lambda p : a \wedge b. \langle \pi_2 p, \pi_1 p \rangle$$

different names for the same theory

simply typed lambda calculus

$\parallel$

simple type theory = STT

$\parallel$

$\lambda \rightarrow$

3 typing rules

$\nVdash$

7 typing rules

same set of well-typed terms

simply typed lambda calculus: syntax and rules

syntax

$A, B ::= a \mid A \rightarrow B$	types
$M, N ::= x \mid MN \mid \lambda x : A. M$	preterms
$\Gamma ::= \cdot \mid \Gamma, x : A$	contexts
$\mathcal{J} ::= \Gamma \vdash M : A$	judgments

rules

$$\frac{}{\Gamma \vdash x : A} \quad \text{for } (x : A) \in \Gamma$$
$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

well-typedness

$\Gamma \vdash M : A$ is derivable $\implies M$ is well-typed

well-typed preterms are called **terms**

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. f x$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

$$\Gamma \vdash M : A$$

example type derivation

$\lambda f : a \rightarrow b. \lambda x : a. fx$

$\frac{\cdot \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}{\Gamma \vdash M : A}$

$\Gamma \vdash M : A$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. f x$$

$$\cdot \vdash (\lambda f : a \rightarrow b. \lambda x : a. f x) : (a \rightarrow b) \rightarrow a \rightarrow b$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\cdot \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\cdot \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\cdot \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$\lambda f : a \rightarrow b. \lambda x : a. fx$

$\frac{}{\Gamma \vdash \cdot : ()}$
 $\frac{}{\Gamma \vdash x : A}$
 $\frac{}{\Gamma \vdash f : a \rightarrow b}$
 $\frac{}{\Gamma \vdash (a \rightarrow b) \rightarrow a \rightarrow b}$

$\frac{}{\Gamma \vdash x : A}$
 $\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B}$
 $\frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\begin{array}{c} \Gamma \quad x \quad A \quad M \quad B \\ \hline \cdot, f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b \end{array}}{\begin{array}{c} \Gamma \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b \end{array}}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\cdot, f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}{\cdot \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\frac{f : a \rightarrow b, x : a \vdash fx : b}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\frac{f : a \rightarrow b, x : a \vdash \textcolor{red}{f}x : b}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash \textcolor{red}{F}M : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\frac{\frac{\Gamma}{f : a \rightarrow b, x : a \vdash \boxed{f} : a \rightarrow b} \quad \frac{\frac{F \quad A \quad B}{f : a \rightarrow b, x : a \vdash \boxed{f} : a \rightarrow b}}{\frac{f : a \rightarrow b, x : a \vdash fx : b}} \quad \frac{\frac{\Gamma}{f : a \rightarrow b, x : a \vdash \boxed{x} : a} \quad \frac{M \quad A}{f : a \rightarrow b, x : a \vdash \boxed{x} : a}}{\frac{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\frac{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\frac{\frac{f : a \rightarrow b, x : a \vdash f : a \rightarrow b \quad f : a \rightarrow b, x : a \vdash x : a}{f : a \rightarrow b, x : a \vdash fx : b}}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\frac{\frac{f : a \rightarrow b, x : a \vdash f : a \rightarrow b}{f : a \rightarrow b, x : a \vdash fx : b}}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$$\lambda f : a \rightarrow b. \lambda x : a. fx$$

$$\frac{\frac{\frac{f : a \rightarrow b, x : a \vdash f : a \rightarrow b}{f : a \rightarrow b, x : a \vdash fx : b}}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

example type derivation

$\lambda f : a \rightarrow b. \lambda x : a. fx$

$$\frac{\frac{\frac{f : a \rightarrow b, x : a \vdash f : a \rightarrow b}{f : a \rightarrow b, x : a \vdash fx : b}}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

$$\frac{}{\Gamma \vdash x : A} \quad \frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash (\lambda x : A. M) : A \rightarrow B} \quad \frac{\Gamma \vdash F : A \rightarrow B \quad \Gamma \vdash M : A}{\Gamma \vdash FM : B}$$

proofs versus type derivations: type derivation

$$\frac{\frac{\frac{}{f : a \rightarrow b, x : a \vdash f : a \rightarrow b} \quad \frac{}{f : a \rightarrow b, x : a \vdash x : a}}{f : a \rightarrow b, x : a \vdash fx : b}}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

proofs versus type derivations: type derivation

$$\frac{\frac{\frac{}{f : a \rightarrow b, x : a \vdash f : a \rightarrow b} \quad \frac{}{f : a \rightarrow b, x : a \vdash x : a}}{f : a \rightarrow b, x : a \vdash fx : b}}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b}}{\vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b}$$

proofs versus type derivations: type derivation

$$\begin{array}{c}
 \overline{f : a \rightarrow b, x : a \vdash f : a \rightarrow b} \quad \overline{f : a \rightarrow b, x : a \vdash x : a} \\
 \hline
 f : a \rightarrow b, x : a \vdash fx : b \\
 \hline
 f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b \\
 \hline
 \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b
 \end{array}$$

$$\begin{array}{c}
 \overline{a \rightarrow b, a \vdash a \rightarrow b} \quad \overline{a \rightarrow b, a \vdash a} \\
 \hline
 a \rightarrow b, a \vdash b \\
 \hline
 a \rightarrow b \vdash a \rightarrow b \\
 \hline
 \vdash (a \rightarrow b) \rightarrow a \rightarrow b
 \end{array}$$

proofs versus type derivations: type derivation

$$\begin{array}{c}
 \frac{\frac{}{f : a \rightarrow b, x : a \vdash f : a \rightarrow b} \quad \frac{}{f : a \rightarrow b, x : a \vdash x : a}}{\frac{}{f : a \rightarrow b, x : a \vdash fx : b}} \\
 \frac{}{f : a \rightarrow b \vdash (\lambda x : a. fx) : a \rightarrow b} \\
 \hline
 \vdash (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b
 \end{array}$$

$$\begin{array}{c}
 \frac{}{a \rightarrow b, a \vdash a \rightarrow b} \text{ass} \quad \frac{}{a \rightarrow b, a \vdash a} \text{ass} \\
 \hline
 \frac{}{a \rightarrow b, a \vdash b} E \rightarrow \\
 \frac{}{a \rightarrow b, a \vdash b} I \rightarrow \\
 \frac{}{a \rightarrow b \vdash a \rightarrow b} I \rightarrow \\
 \hline
 \vdash (a \rightarrow b) \rightarrow a \rightarrow b
 \end{array}$$

proofs versus type derivations: proof

$$\begin{array}{c}
 \frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow \\
 \frac{a \rightarrow b}{a \rightarrow b} I[x] \rightarrow \\
 \frac{}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow
 \end{array}$$

$$\begin{array}{c}
 \frac{}{a \rightarrow b, a \vdash a \rightarrow b} \text{ass} \quad \frac{}{a \rightarrow b, a \vdash a} \text{ass} \\
 \frac{}{a \rightarrow b, a \vdash b} E \rightarrow \\
 \frac{a \rightarrow b, a \vdash b}{a \rightarrow b \vdash a \rightarrow b} I \rightarrow \\
 \frac{a \rightarrow b \vdash a \rightarrow b}{\vdash (a \rightarrow b) \rightarrow a \rightarrow b} I \rightarrow
 \end{array}$$

proofs versus type derivations: proof term

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

$$\lambda f : a \rightarrow b. \lambda x : a. f x$$

Curry-Howard correspondence

propositions

types

implication $\longleftrightarrow$ function type

proof rules

proof terms

introduction rule $\longleftrightarrow$ lambda abstraction

elimination rule $\longleftrightarrow$ function application

assumption rule $\longleftrightarrow$ variable

how to read proof terms

$$M := (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

how to read proof terms

$$M := (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

this proof term M is a function
that takes as its first argument a proof of $a \rightarrow b$ called f
and as its second argument a proof of a called x
and then maps those to a proof of b

how to read proof terms

$$M := (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

this proof term M is a function
that takes as its first argument a proof of $a \rightarrow b$ called f
and as its second argument a proof of a called x
and then maps those to a proof of b

the argument f is itself a function that
maps proofs of a to proofs of b
(conform the BHK-interpretation)

x is an inhabitant of the type a
which corresponds to the proposition a

how to read proof terms

$$M := (\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

this proof term M is a function
that takes as its first argument a proof of $a \rightarrow b$ called f
and as its second argument a proof of a called x
and then maps those to a proof of b

the argument f is itself a function that
maps proofs of a to proofs of b
(conform the BHK-interpretation)

x is an inhabitant of the type a
which corresponds to the proposition a

to get a proof of b , the function f is applied to x
and the result of that application then is the result of M

Rocq

the example

Rocq

the example

```
Parameter a b : Prop.      a is declared  
                           b is declared
```

the example

Parameter a b : Prop.

1 subgoal (ID 1)

Lemma one :

(a -> b) -> a -> b.

=====

(a -> b) -> a -> b

$(a \rightarrow b) \rightarrow a \rightarrow b$

Rocq

the example

Parameter a b : Prop.

1 subgoal (ID 2)

Lemma one :

f : a -> b

(a -> b) -> a -> b.

=====

intro f.

a -> b

$$\frac{a \rightarrow b}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

Rocq

the example

Parameter a b : Prop.

1 subgoal (ID 3)

Lemma one :

f : a -> b

(a -> b) -> a -> b.

x : a

intro f.

=====

intro x.

b

$$\frac{\frac{b}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

Rocq

the example

Parameter a b : Prop.

1 subgoal (ID 4)

Lemma one :

f : a -> b

(a -> b) -> a -> b.

x : a

intro f.

=====

intro x.

a

apply f.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad a}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

the example

Parameter a b : Prop.

No more subgoals.

Lemma one :

(a -> b) -> a -> b.

intro f.

intro x.

apply f.

exact x.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

the example

Parameter a b : Prop.

Lemma one :

```
(a -> b) -> a -> b.
intro f.
intro x.
apply f.
exact x.
Qed.
```

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

the example

Parameter a b : Prop.

one

: (a -> b) -> a -> b

Lemma one :

(a -> b) -> a -> b.

intro f.

intro x.

apply f.

exact x.

Qed.

Check one.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

Rocq

the example

Parameter a b : Prop.

one = fun (f : a -> b) (x : a) => f x
: (a -> b) -> a -> b

Lemma one :

(a -> b) -> a -> b.
intro f.
intro x.
apply f.
exact x.
Qed.

Arguments one _%function_scope

Check one.

Print one.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

$$(\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

Rocq

the example

Parameter a b : Prop.

1 subgoal (ID 3)

Lemma one :

f : a -> b

(a -> b) -> a -> b.

x : a

intro f.

=====

intro x.

b

apply f.

exact x.

Qed.

Check one.

Print one.

$$\begin{array}{c}
 \frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow \\
 \frac{b}{a \rightarrow b} I[x] \rightarrow \\
 \frac{a \rightarrow b}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow
 \end{array}$$

$$(\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

Rocq

the example

Parameter a b : Prop.

1 subgoal (ID 3)

Lemma one :

H : a -> b

(a -> b) -> a -> b.

H0 : a

intros.

=====

apply H.

b

exact H0.

Qed.

Check one.

Print one.

$$\frac{\frac{\frac{[a \rightarrow b^H] \quad [a^{H_0}]}{b} E \rightarrow}{a \rightarrow b} I[H_0] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[H] \rightarrow$$

$$(\lambda H : a \rightarrow b. \lambda H_0 : a. H H_0) : (a \rightarrow b) \rightarrow a \rightarrow b$$

Rocq

the example

Parameter a b : Prop.

1 subgoal (ID 3)

Lemma one :

f : a -> b

(a -> b) -> a -> b.

x : a

intros f x.

=====

apply f.

b

exact x.

Qed.

Check one.

Print one.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

$$(\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

the example

Parameter a b : Prop. No more subgoals.

Lemma one :

(a -> b) -> a -> b.

intros f x.

apply f.

exact x.

Qed.

Check one.

Print one.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

$$(\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

the example

Parameter a b : Prop. No more subgoals.

Lemma one :

 (a -> b) -> a -> b.

intros f x.

apply f.

apply x.

Qed.

Check one.

Print one.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

$$(\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

the example

Parameter a b : Prop. No more subgoals.

Lemma one :

 (a -> b) -> a -> b.

auto.

Qed.

Check one.

Print one.

Rocq

the example

Parameter a b : Prop.

one = fun H : a → b => H
 : (a → b) → a → b

Lemma one :

 (a → b) → a → b.
auto.
Qed.

Arguments one _%function_scope

Check one.

Print one.

$$\frac{[a \rightarrow b^H]}{(a \rightarrow b) \rightarrow a \rightarrow b} I[H] \rightarrow$$

$$(\lambda H : a \rightarrow b. H) : (a \rightarrow b) \rightarrow a \rightarrow b$$

Rocq

the example

Parameter a b : Prop.

one = fun (f : a -> b) (x : a) => f x
 : (a -> b) -> a -> b

Lemma one :

 (a -> b) -> a -> b.

intros f x.

apply f.

apply x.

Qed.

Check one.

Print one.

$$\frac{\frac{\frac{[a \rightarrow b^f] \quad [a^x]}{b} E \rightarrow}{a \rightarrow b} I[x] \rightarrow}{(a \rightarrow b) \rightarrow a \rightarrow b} I[f] \rightarrow$$

$$(\lambda f : a \rightarrow b. \lambda x : a. fx) : (a \rightarrow b) \rightarrow a \rightarrow b$$

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 2)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

=====

a $\vee$ b $\rightarrow$ b $\vee$ a

$$a \vee b \rightarrow b \vee a$$

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 3)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

x : a $\vee$ b

=====

b $\vee$ a

$$\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow$$

example with disjunction

Parameter a b : Prop.

2 subgoals (ID 4)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

x : a $\vee$ b

=====

a $\rightarrow$ b $\vee$ a

subgoal 2 (ID 5) is:

b $\rightarrow$ b $\vee$ a

$$\frac{\begin{array}{ccc} [a \vee b^x] & a \rightarrow b \vee a & b \rightarrow b \vee a \end{array}}{\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow} E\vee$$

example with disjunction

Parameter a b : Prop.

2 subgoals (ID 6)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

x : a $\vee$ b

y : a

=====

b $\vee$ a

subgoal 2 (ID 5) is:

b $\rightarrow$ b $\vee$ a

$$\frac{[a \vee b^x] \quad \frac{b \vee a}{a \rightarrow b \vee a} I[y] \rightarrow \quad b \rightarrow b \vee a}{\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow} E\vee$$

example with disjunction

Parameter a b : Prop.

2 subgoals (ID 8)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

right.

x : a $\vee$ b

y : a

=====

a

subgoal 2 (ID 5) is:

b $\rightarrow$ b $\vee$ a

$$\frac{\frac{[a \vee b^x] \quad \frac{\frac{a}{b \vee a} \text{Ir}\vee \quad a \rightarrow b \vee a}{I[y] \rightarrow} \quad b \rightarrow b \vee a}{b \vee a} E\vee}{\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow}$$

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 5)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

right.

apply y.

x : a $\vee$ b

=====

b $\rightarrow$ b $\vee$ a

$$\frac{\frac{\frac{[a \vee b^x]}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{\frac{[a^y]}{b \vee a} Ir \vee \quad b \rightarrow b \vee a}{a \vee b \rightarrow b \vee a} E \vee}{a \vee b \rightarrow b \vee a} I[x] \rightarrow$$

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 9)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

right.

apply y.

intro z.

x : a $\vee$ b

z : b

b $\vee$ a

$$\begin{array}{c}
 \frac{[a \vee b^x] \quad \frac{\frac{[a^y]}{b \vee a} Ir\vee \quad \frac{b \vee a}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{b \vee a}{b \rightarrow b \vee a} I[z] \rightarrow}{E\vee}}{a \vee b \rightarrow b \vee a} I[x] \rightarrow
 \end{array}$$

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 11)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

right.

apply y.

intro z.

left.

x : a $\vee$ b

z : b

=====

b

$$\frac{\frac{[a \vee b^x]}{a \rightarrow b \vee a} \frac{\frac{[a^y]}{b \vee a} Ir\vee}{I[y] \rightarrow} \quad \frac{\frac{b}{b \vee a} Il\vee}{b \rightarrow b \vee a} I[z] \rightarrow}{\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow} E\vee$$

example with disjunction

Parameter a b : Prop.

No more subgoals.

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

right.

apply y.

intro z.

left.

apply z.

$$\frac{\frac{[a \vee b^x]}{a \rightarrow b \vee a} \quad \frac{\frac{\frac{[a^y]}{b \vee a} Ir\vee}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{\frac{\frac{[b^z]}{b \vee a} Il\vee}{b \rightarrow b \vee a} I[z] \rightarrow}{b \vee a} E\vee}{a \vee b \rightarrow b \vee a} I[x] \rightarrow$$

example with disjunction

Parameter a b : Prop.

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

intro x.

elim x.

intro y.

right.

apply y.

intro z.

left.

apply z.

Qed.

$$\frac{[a \vee b^x] \quad \frac{\frac{[a^y]}{b \vee a} Ir\vee \quad \frac{[b^z]}{b \vee a} Il\vee}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{b \rightarrow b \vee a}{I[z] \rightarrow} E\vee}{\frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow} EV$$

example with disjunction

Parameter a b : Prop.

2 subgoals (ID 4)

Lemma two :

$x : a \vee b$

$a \vee b \rightarrow b \vee a$.

=====

intro x.

$a \rightarrow b \vee a$

elim x.

intro y.

subgoal 2 (ID 5) is:

right.

$b \rightarrow b \vee a$

apply y.

intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

2 subgoals (ID 4)

Lemma two :

x : a $\vee$ b

a $\vee$ b $\rightarrow$ b $\vee$ a.

=====

intro x.

a $\rightarrow$ b $\vee$ a

elim x.

- intro y.

subgoal 2 (ID 5) is:

right.

b $\rightarrow$ b $\vee$ a

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 4)

Lemma two :

x : a $\vee$ b

a $\vee$ b $\rightarrow$ b $\vee$ a.

=====

intro x.

a $\rightarrow$ b $\vee$ a

elim x.

- intro y.

right.

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 6)

Lemma two :

x : a $\vee$ b

a $\vee$ b $\rightarrow$ b $\vee$ a.

y : a

intro x.

=====

elim x.

b $\vee$ a

- intro y.

right.

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 8)

Lemma two :

x : a $\vee$ b

a $\vee$ b $\rightarrow$ b $\vee$ a.

y : a

intro x.

=====

elim x.

a

- intro y.

right.

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 5)

Lemma two :

subgoal 1 (ID 5) is:

a $\vee$ b $\rightarrow$ b $\vee$ a.

b $\rightarrow$ b $\vee$ a

intro x.

elim x.

- intro y.

right.

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 5)

Lemma two :

x : a $\vee$ b

a $\vee$ b $\rightarrow$ b $\vee$ a.

=====

intro x.

b $\rightarrow$ b $\vee$ a

elim x.

- intro y.

right.

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 6)

Lemma two :

x : a $\vee$ b

a $\vee$ b $\rightarrow$ b $\vee$ a.

y : a

intro x.

=====

elim x.

b $\vee$ a

- intro y.

right.

apply y.

- intro z.

left.

apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 9)

Lemma two :

y : a

a $\vee$ b $\rightarrow$ b $\vee$ a.

=====

intro x.

b $\vee$ a

destruct x as [y|z].

- right.

 apply y.

- left.

 apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 9)

Lemma two :

y : a

a $\vee$ b $\rightarrow$ b $\vee$ a.

=====

intro x.

b $\vee$ a

destruct x as [y|z].

- right.

 apply y.

- left.

 apply z.

Qed.

example with disjunction

Parameter a b : Prop.

1 subgoal (ID 8)

Lemma two :

a $\vee$ b $\rightarrow$ b $\vee$ a.

y : a

=====

intros [y|z].

b $\vee$ a

- right.

 apply y.

- left.

 apply z.

Qed.

example with disjunction

Parameter a b : Prop.

Lemma two :

$a \vee b \rightarrow b \vee a$.

intros [y|z].

- right.

 apply y.

- left.

 apply z.

Qed.

$$\frac{\frac{[a^y]}{b \vee a} Ir\vee \quad \frac{[b^z]}{b \vee a} Il\vee}{\frac{[a \vee b^x]}{a \rightarrow b \vee a} I[y] \rightarrow \quad \frac{b \rightarrow b \vee a}{I[z] \rightarrow} E\vee} \frac{b \vee a}{a \vee b \rightarrow b \vee a} I[x] \rightarrow$$

example with conjunction

Parameter a b : Prop.

1 subgoal (ID 3)

Lemma three :

a /\ b -> b /\ a.

=====

a /\ b -> b /\ a

$$a \wedge b \rightarrow b \wedge a$$

example with conjunction

Parameter a b : Prop.

1 subgoal (ID 8)

Lemma three :

a /\ b -> b /\ a.

intros [y z].

y : a

z : b

=====

b /\ a

$$\frac{\frac{\frac{b \wedge a}{b \rightarrow b \wedge a} I[z] \rightarrow}{a \rightarrow b \rightarrow b \wedge a} I[y] \rightarrow}{\frac{[a \wedge b^x] \quad a \rightarrow b \rightarrow b \wedge a}{b \wedge a} E \wedge} I[x] \rightarrow$$

$a \wedge b \rightarrow b \wedge a$

example with conjunction

Parameter a b : Prop.

2 subgoals (ID 10)

Lemma three :

a /\ b -> b /\ a.

intros [y z].

split.

y : a

z : b

=====

b

subgoal 2 (ID 11) is:

a

$$\frac{\frac{\frac{b \quad a}{b \wedge a} I\wedge \quad \frac{b \rightarrow b \wedge a}{a \rightarrow b \rightarrow b \wedge a} I[z] \rightarrow \quad \frac{I[y] \rightarrow}{E\wedge}}{[a \wedge b^x] \quad \frac{b \wedge a}{a \wedge b \rightarrow b \wedge a} I[x] \rightarrow}$$

example with conjunction

Parameter a b : Prop.

1 subgoal (ID 10)

Lemma three :

a /\ b -> b /\ a.

intros [y z].

split.

-

y : a

z : b

=====

b

$$\frac{\frac{\frac{b}{I\wedge} \quad \frac{a}{I\wedge}}{b \wedge a} \quad \frac{I[z] \rightarrow}{b \rightarrow b \wedge a} \quad \frac{I[y] \rightarrow}{a \rightarrow b \rightarrow b \wedge a}}{[a \wedge b^x] \quad E\wedge} \quad \frac{b \wedge a}{a \wedge b \rightarrow b \wedge a} I[x] \rightarrow$$

example with conjunction

Parameter a b : Prop.

1 subgoal (ID 11)

Lemma three :

subgoal 1 (ID 11) is:

a /\ b -> b /\ a.

a

intros [y z].

split.

- apply z.

$$\frac{\frac{\frac{[a \wedge b^x]}{a \rightarrow b \rightarrow b \wedge a} E \wedge \quad \frac{\frac{\frac{[b^z] \quad a}{b \wedge a} I \wedge \quad \frac{b \rightarrow b \wedge a}{I[z] \rightarrow} \quad \frac{a \rightarrow b \rightarrow b \wedge a}{I[y] \rightarrow}}{b \wedge a} I[x] \rightarrow}{a \wedge b \rightarrow b \wedge a} I[x] \rightarrow$$

example with conjunction

Parameter a b : Prop.

1 subgoal (ID 11)

Lemma three :

a /\ b -> b /\ a.

intros [y z].

split.

- apply z.

-

y : a

z : b

=====

a

$$\frac{\frac{\frac{[a \wedge b^x]}{a \rightarrow b \rightarrow b \wedge a} E \wedge \quad \frac{\frac{\frac{[b^z]}{b \wedge a} I \wedge \quad \frac{a}{b \rightarrow b \wedge a} I[z] \rightarrow}{a \rightarrow b \rightarrow b \wedge a} I[y] \rightarrow}{b \wedge a} I[x] \rightarrow}{a \wedge b \rightarrow b \wedge a}$$

example with conjunction

Parameter a b : Prop.

No more subgoals.

Lemma three :

a /\ b -> b /\ a.

intros [y z].

split.

- apply z.

- apply y.

$$\frac{\frac{\frac{[b^z] \quad [a^y]}{b \wedge a} I \wedge}{b \rightarrow b \wedge a} I[z] \rightarrow}{a \rightarrow b \rightarrow b \wedge a} I[y] \rightarrow}{[a \wedge b^x] \quad a \wedge b \rightarrow b \wedge a} E \wedge$$

example with conjunction

Parameter a b : Prop.

Lemma three :

a /\ b -> b /\ a.

intros [y z].

split.

- apply z.

- apply y.

Qed.

$$\frac{\frac{\frac{[b^z] \quad [a^y]}{b \wedge a} I \wedge \quad \frac{b \wedge a}{b \rightarrow b \wedge a} I[z] \rightarrow \quad \frac{a \rightarrow b \rightarrow b \wedge a}{a \rightarrow b \rightarrow b \wedge a} I[y] \rightarrow}{[a \wedge b^x] \quad a \rightarrow b \rightarrow b \wedge a} E \wedge}{b \wedge a} \frac{b \wedge a}{a \wedge b \rightarrow b \wedge a} I[x] \rightarrow$$

example with conjunction

Parameter a b : Prop.

Lemma three :

$a \wedge b \rightarrow b \wedge a$.

intros [y z].

split.

- apply z.

- apply y.

Qed.

$$\frac{\frac{\frac{[a \wedge b^x] \quad \frac{\frac{[b^z] \quad [a^y]}{b \wedge a} I \wedge}{b \rightarrow b \wedge a} I[z] \rightarrow}{a \rightarrow b \rightarrow b \wedge a} I[y] \rightarrow}{b \wedge a} E \wedge}{a \wedge b \rightarrow b \wedge a} I[x] \rightarrow$$

tactics for proof rules

$I \rightarrow I \neg$	intro intros
$E \rightarrow E \neg$	apply
ass	exact apply
$E \wedge E \vee E \perp$	elim destruct intros with pattern
$I \wedge$	split
$Il \vee$	left
$Ir \vee$	right
$I \top$	apply I

tactics for proof rules

$I \rightarrow I \neg$	intro intros
$E \rightarrow E \neg$	apply
ass	exact apply
$E \wedge E \vee E \perp$	elim destruct intros with pattern
$I \wedge$	split
$Il \vee$	left
$Ir \vee$	right
$I \top$	apply I I : True

bullets

structure tactic scripts according to subgoals

related bullets need to match:

- -- --- ---- etc.

+ ++ +++ ++++ etc.

* ** *** **** etc.

intro patterns

only works with `intros`, not with `intro`
the pattern needs to match the shape of assumptions

goal	tactic
$A \rightarrow G$	<code>intros HA.</code>
$A \rightarrow B \rightarrow G$	<code>intros HA HB.</code>
$A \wedge B \rightarrow G$	<code>intros [HA HB].</code>
$A \vee B \rightarrow G$	<code>intros [HA HB].</code>
$A \vee (B \wedge C) \rightarrow D \rightarrow G$	<code>intros [HA [HB HC]] HD.</code>

apply

$$H : A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B$$

goal B $\xrightarrow{\text{apply } H}$ new goals $A_1, \dots, A_n$

apply

the number n of antecedents in the type of H may be zero
 H may be an arbitrary term

$$H : A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B$$

goal B $\xrightarrow{\text{apply } H}$ new goals $A_1, \dots, A_n$

apply

the number n of antecedents in the type of H may be zero
 H may be an arbitrary term

$$H : A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B$$

goal B $\xrightarrow{\text{apply } H}$ new goals $A_1, \dots, A_n$

$$\begin{array}{c} \frac{[A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B^H] \quad A_1}{A_2 \rightarrow \cdots \rightarrow A_n \rightarrow B} E \rightarrow \\ \hline \frac{A_2 \rightarrow \cdots \rightarrow A_n \rightarrow B \quad A_2}{A_3 \rightarrow \cdots \rightarrow A_n \rightarrow B} E \rightarrow \\ \vdots \\ \frac{A_{n-1} \rightarrow A_n \rightarrow B \quad A_{n-1}}{A_n \rightarrow B} E \rightarrow \\ \hline \frac{A_n \rightarrow B \quad A_n}{B} E \rightarrow \end{array}$$

elim

$$H : A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B \otimes C$$

elim H and destruct eliminate the connective $\otimes$
but also generate extra subgoals $A_1, \dots, A_n$

elim

$$H : A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B \otimes C$$

elim H and destruct eliminate the connective $\otimes$
 but also generate extra subgoals $A_1, \dots, A_n$

$$\begin{array}{c}
 \frac{[A_1 \rightarrow \cdots \rightarrow A_n \rightarrow B \otimes C^H] \quad A_1}{A_2 \rightarrow \cdots \rightarrow A_n \rightarrow B \otimes C} E \rightarrow \\
 \hline
 \vdots \\
 \frac{A_{n-1} \rightarrow A_n \rightarrow B \otimes C \quad A_{n-1}}{A_n \rightarrow B \otimes C} E \rightarrow \\
 \hline
 \frac{A_n \rightarrow B \otimes C \quad A_n}{B \otimes C} E \rightarrow \\
 \hline
 \frac{B \otimes C \quad \cdots}{\dots} E \otimes
 \end{array}$$

summary

minimal logic	simple type theory	Rocq
only implication		
introduction rule	lambda abstraction	intro
elimination rule	function application	apply

summary

full constructive logic

Rocq type theory

Rocq

implication

introduction rule

lambda abstraction

`intro`

elimination rule

function application

`apply`

the other connectives

introduction rules

(in three weeks)

constructor
left/right

elimination rules

(in three weeks)

`elim`

summary

full constructive logic

Rocq type theory

Rocq

implication

introduction rule
elimination rule

lambda abstraction
function application

intro
apply

the other connectives

introduction rules

(in three weeks)

constructor
left/right

elimination rules

(in three weeks)

elim

for more about the tactics
read the Rocq manual!



T-shirt

a lambda calculus self-interpreter

$$U_k^i \equiv \lambda x_1 \dots x_k. x_i$$
$$\langle M_1, \dots, M_k \rangle \equiv \lambda z. z M_1 \dots M_k$$

$$\langle \langle K, S, C \rangle \rangle \ulcorner M \urcorner \twoheadrightarrow M$$

$$\ulcorner x \urcorner \equiv \lambda e. e U_3^1 x e$$
$$\ulcorner P Q \urcorner \equiv \lambda e. e U_3^2 \ulcorner P \urcorner \ulcorner Q \urcorner e$$
$$\ulcorner \lambda x. P \urcorner \equiv \lambda e. e U_3^3 (\lambda x. \ulcorner P \urcorner) e$$

$$K \equiv \lambda x y. x$$

$$S \equiv \lambda x y z. x z (y z)$$

$$C \equiv \lambda x y z. x z y$$

table of contents

contents

type systems and logics

propositional logic

simply typed lambda calculus

Rocq

T-shirt

table of contents